

Serial No. 10/008,620

Page 2

LIST OF CLAIMS, SHOWING THE STATUS OF EACH CLAIM

Underlining denotes added text while bracketing denotes deleted text.

IN THE CLAIMS:

Please cancel Claims 7, 17, as indicated in the following list.

1. A method of producing a library of mutant nucleic acid molecules comprising:
 - (a) obtaining a template nucleic acid;
 - (b) preparing a first oligonucleotide corresponding to a first desired mutation within said template nucleic acid;
 - (c) preparing a second oligonucleotide corresponding to a second desired mutation within said template nucleic acid;
 - (d) mixing the oligonucleotides prepared in said steps (b) and (c) so as to hybridize said oligonucleotides to said template nucleic acid;
 - (e) subjecting the mixture of step (d) to the linear cyclic amplification reaction to produce a library of mutant template nucleic acids.
2. The method according to claim 1, wherein said oligonucleotides in said steps (b) and (c) are discontiguous.
3. The method according to claim 1, wherein said step first and second oligonucleotides are present in less than saturation concentration.
4. The method according to claim 1, wherein the mixture of said step (d) further comprises non-mutagenic oligonucleotides corresponding to either or both of said first and second oligonucleotides.
5. The method according to claim 1, wherein said template nucleic acid corresponds to a desired protein product.
6. The method according to claim 4, wherein said protein product comprises an enzyme, hormone, vaccine, peptide therapeutic or antibody.

Serial No. 10/008,620

Page 3

7. (canceled)
8. A method for producing a library of mutant nucleic acid molecules comprising the steps of:
- (a) obtaining a template nucleic acid;
 - (b) preparing two or more primers corresponding to the template nucleic acid, wherein at least one primer is in opposite orientation to the remaining primers and at least one primer is a mutagenic primer corresponding to a desired mutation;
 - (c) mixing the primers in said step (b) so as to hybridize said primers to said template nucleic acid; and
 - (d) subjecting the mixture of step (c) to the linear cyclic amplification reaction to produce a library of mutant template nucleic acids.
9. The method of claim 8, wherein said two or more primers comprises 3 to 15 primers or 4 to 7 primers.
10. The method of claim 8, wherein said primers in said step (b) are discontiguous.
11. The method according to claim 8, wherein said primers in step (b) are present in less than saturation concentration.
12. The method of claim 8, wherein all said primers in step (b) are mutagenic primers.
13. The method of claim 8, wherein said at least one mutagenic primer comprises 1 to 12 nucleotide mutations.
14. The method of claim 8, wherein said at least one mutagenic primer encodes 1 to 4 amino acid mutations.
15. The method according to claim 8, wherein said template nucleic acid corresponds to a desired protein product.

Serial No. 10/008,620

Page 4

B
Canc
16. The method according to claim 15, wherein said protein product comprises an enzyme, hormone, vaccine, peptide therapeutic or antibody.

17. (canceled)